

Tutorial for Introductory Analysis of Daily Precipitation Data with hydroTSM

Mauricio Zambrano-Bigiarini

Last Update: 30-May-2013

1 Installation

Installing the latest stable version (from CRAN):

```
> install.packages("hydroTSM")
```

Alternatively, you can also try the under-development version (from rforge):

```
> install.packages("hydroTSM", , "http://rforge.net/", type="source")
```

2 Setting Up the Environment

1. Loading the *hydroTSM* library, which contains data and functions used in this analysis.

```
> library(hydroTSM)
```

2. Loading daily precipitation data at the station San Martino di Castrozza, Trento Province, Italy, with data from 01/Jan/1921 to 31/Dec/1990.

```
> data(SanMartinoPPts)
```

3. Selecting only a 6-years time slice for the analysis

```
> x <- window(SanMartinoPPts, start=as.Date("1985-01-01"))
```

4. Monthly values of precipitation

```
> ( m <- daily2monthly(x, FUN=sum) )
```

1985-01-01	1985-02-01	1985-03-01	1985-04-01	1985-05-01	1985-06-01	1985-07-01	
141.2	7.0	140.6	72.0	175.6	131.4	85.4	
159.4	27.2	58.4	101.8	54.8	75.8	131.6	
59.6	237.8	108.2	144.8	81.2	141.0	69.8	
38.2	44.4	20.4	46.8	111.0	45.6	98.4	
212.0	153.8	221.8	175.0	90.6	278.8	164.8	
29.8	118.0	49.8	22.4	100.6	187.4	193.0	
120.4	149.2	61.2	136.4	10.0	59.4	0.0	
152.6	46.2	365.4	77.4	241.6	302.8	114.4	
65.4	12.8	145.0	110.6	51.6	12.4	65.8	
127.0	74.4	175.0	143.8	90.8	106.0	153.0	
326.6	106.0						

5. Dates of the daily values of 'x'

```
> dates <- time(x)
```

6. Amount of years in 'x' (needed for computations)

```
> ( nyears <- yip(from=start(x), to=end(x), out.type="nmbr" ) )
```

```
[1] 6
```

3 Basic Exploratory Data Analysis


- Summary statistics

```
> smry(x)
```

	Index	x
Min.	1985-01-01	0.0000
1st Qu.	1986-07-02	0.0000
Median	1988-01-01	0.0000
Mean	1988-01-01	3.7470
3rd Qu.	1989-07-01	2.6000
Max.	1990-12-31	122.0000
IQR	<NA>	2.6000
sd	<NA>	10.0428
cv	<NA>	2.6800
Skewness	<NA>	5.3512
Kurtosis	<NA>	39.1619
NA's	<NA>	0.0000
n	<NA>	2191.0000

- Using the *hydroplot* function, which (by default) plots 9 different graphs: 3 ts plots, 3 boxplots and 3 histograms summarizing 'x'. For this example, only daily and monthly plots are produced, and only data starting on 01-Jan-1987 are plotted.

```
> hydroplot(x, var.type="Precipitation", main="at San Martino",
+           pfreq = "dm", from="1987-01-01")
```



3. Amount of days with information (not NA) per year

```
> dwi(x)
```

```
1985 1986 1987 1988 1989 1990  
365 365 365 366 365 365
```

4. Amount of days with information (not NA) per month per year


```
> dwi(x, out.unit="mpy")
```

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1985	31	28	31	30	31	30	31	31	30	31	30	31
1986	31	28	31	30	31	30	31	31	30	31	30	31
1987	31	28	31	30	31	30	31	31	30	31	30	31
1988	31	29	31	30	31	30	31	31	30	31	30	31
1989	31	28	31	30	31	30	31	31	30	31	30	31
1990	31	28	31	30	31	30	31	31	30	31	30	31

5. Plotting the monthly precipitation values for each year, useful for identifying dry/wet months.

```
> # Daily zoo to monthly zoo  
> m <- daily2monthly(x, FUN=sum, na.rm=TRUE)  
> # Creating a matrix with monthly values per year in each column  
> M <- matrix(m, ncol=12, byrow=TRUE)  
> colnames(M) <- month.abb  
> rownames(M) <- unique(format(time(m), "%Y"))  
> # Plotting the monthly precipitation values  
> require(lattice)  
> print(matrixplot(M, ColorRamp="Precipitation",  
+ main="Monthly precipitation at San Martino st., [mm/month]"))
```

Monthly precipitation at San Martino st., [mm/month]



4 Annual Analysis

1. Annual values of precipitation

```
> daily2annual(x, FUN=sum, na.rm=TRUE)  
  
1985-01-01 1986-01-01 1987-01-01 1988-01-01 1989-01-01 1990-01-01  
1154.8     1152.8     1628.4     1207.8     1634.2     1432.4
```

2. Average annual precipitation

Obvious way:

```
> mean(daily2annual(x, FUN=sum, na.rm=TRUE))  
  
[1] 1368.4
```

Another way (more useful for streamflows, where `FUN=mean`):

The function `annualfunction` applies `FUN` twice over `x`: (i) firstly, over all the elements of `x` belonging to the same year, in order to obtain the corresponding annual values, and (ii) secondly, over all the annual values of `x` previously obtained, in order to obtain a single annual value.

```
> annualfunction(x, FUN=sum, na.rm=TRUE) / nyears  
  
value  
1368.4
```

5 Monthly Analysis

1. Median of the monthly values at station 'x'. Not needed, just for looking at these values in the boxplot.

```
> monthlyfunction(m, FUN=median, na.rm=TRUE)  
  
Jan   Feb   Mar   Apr   May   Jun   Jul   Aug   Sep   Oct   Nov   Dec  
63.7  80.4  52.9 113.8 141.9 164.4 132.1 145.1  67.6  97.4 123.4  57.1
```

2. Vector with the three-letter abbreviations for the month names

```
> cmonth <- format(time(m), "%b")
```


3. Creating ordered monthly factors

```
> months <- factor(cmonth, levels=unique(cmonth), ordered=TRUE)
```

4. Boxplot of the monthly values

```
> boxplot(coredata(m) ~ months, col="lightblue", main="Monthly Precipitation",  
+           ylab="Precipitation, [mm]", xlab="Month")
```

Monthly Precipitation



6 Seasonal Analysis

1. Average seasonal values of precipitation

```
> seasonalfunction(x, FUN=sum, na.rm=TRUE) / nyears
```


Season	Average Precipitation [mm]
DJF	213.1333
MAM	369.4000
JJA	470.8000
SON	315.0667

2. Extracting the seasonal values for each year

```
> ( DJF <- dm2seasonal(x, season="DJF", FUN=sum) )  
1985 1986 1987 1988 1989 1990  
148.2 262.2 178.2 197.6 212.0 174.6  
  
> ( MAM <- dm2seasonal(m, season="MAM", FUN=sum) )  
1985 1986 1987 1988 1989 1990  
388.2 405.6 356.0 310.4 489.0 267.2  
  
> ( JJA <- dm2seasonal(m, season="JJA", FUN=sum) )  
1985 1986 1987 1988 1989 1990  
376.2 367.0 550.6 462.6 658.8 409.6  
  
> ( SON <- dm2seasonal(m, season="SON", FUN=sum) )  
1985 1986 1987 1988 1989 1990  
187.4 152.4 534.2 207.6 223.2 585.6
```

3. Plotting the time evolution of the seasonal precipitation values

```
> hydroplot(x, pfreq="seasonal", FUN=sum, stype="default")
```



7 Some Extreme Indices

Common steps for the analysis of this section:

1. Loading daily precipitation data at the station San Martino di Castrozza, Trento Province, Italy, with data from 01/Jan/1921 to 31/Dec/1990.


```
> data(SanMartinoPPts)
```

2. Selecting only a three-year time slice for the analysis

```
> x <- window(SanMartinoPPts, start=as.Date("1988-01-01"))
```

3. Plotting the selected time series

```
> hydroplot(x, ptype="ts", pfreq="o", var.unit="mm")
```



7.1 Heavy Precipitation Days (R10mm)

1. Counting and plotting the number of days in the period where precipitation is > 10 [mm]

```
> ( R10mm <- length( x[x>10] ) )
```

```
[1] 127
```

7.2 Very Wet Days (R95p)

1. Identifying the wet days (daily precipitation ≥ 1 mm):

```
> wet.index <- which(x >= 1)
```

2. Computing the 95th percentile of precipitation on wet days ($PRwn95$):

```
> ( PRwn95 <- quantile(x[wet.index], probs=0.95, na.rm=TRUE) )
```

```
95%
39.75
```

Note 1: this computation was carried out for the three-year time period 1988-1990, not the 30-year period 1961-1990 commonly used.

Note 2: missing values are removed from the computation.

3. Identifying the very wet days (daily precipitation $\geq PRwn95$)

```
> (very.wet.index <- which(x >= PRwn95))
```

```
[1] 30 92 234 287 422 423 461 550 551 674 676 719 939 950 998
[16] 1058 1061 1075
```

4. Computing the total precipitation on the very wet days:

```
> ( R95p <- sum(x[very.wet.index]) )
```


```
[1] 1196.4
```

Note 3: this computation was carried out for the three-year time period 1988-1990, not the 30-year period 1961-1990 commonly used

7.3 5-day Total Precipitation

1. Computing the 5-day total (accumulated) precipitation

```
> x.5max <- rollapply(data=x, width=5, FUN=sum, fill=NA, partial= TRUE,
+ align="center")
> hydroplot(x.5max, ptype="ts+boxplot", pfreq="o", var.unit="mm")
```



2. Maximum annual value of 5-day total precipitation

```
> (x.5max.annual <- daily2annual(x.5max, FUN=max, na.rm=TRUE))
1988-01-01 1989-01-01 1990-01-01
  113.2      170.8      237.2
```

Note 1: for this computation, a moving window centred in the current day is used. If the user wants the 5-day total precipitation accumulated in the 4 days before the current day + the precipitation in the current day, the user have to modify the moving window.

Note 2: For the first two and last two values, the width of the window is adapted to ignore values not within the time series

8 Software Details

This tutorial was built under:

```
[1] "x86_64-unknown-linux-gnu (64-bit)"
[1] "R Under development (unstable) (2013-07-03 r63169)"
[1] "hydroTSM 0.4-1-3"
```